



# First Record of Sea Anemone *Anthopleura handi* Dunn, 1978 (Actiniaria: Actinidae) from Coast across the Indian Mainland

Trideb Kr. Sana <sup>a\*</sup>, Prashant Sharma <sup>a</sup>, Chandran R. <sup>a</sup>  
and Santanu Mitra <sup>b</sup>

<sup>a</sup> General Non Chordata Section, Fireproof Spirit Building, Zoological Survey of India, 27, J. L. Nehru Road, Kolkata, 700016, India.

<sup>b</sup> Crustacea Section, Fireproof Spirit Building, Zoological Survey of India, 27, J. L. Nehru Road, Kolkata, 700016, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.56557/UPJOZ/2024/v45i33880

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.mbmph.com/review-history/3173>

Short Research Article

Received: 25/11/2023

Accepted: 29/01/2024

Published: 30/01/2024

## ABSTRACT

*Anthopleura handi* Dunn, 1978 was previously recorded from Andaman Islands. Here we report *Anthopleura handi* Dunn, 1978 for the first time from coast across the Indian mainland. The sea anemone specimen was collected from intertidal sandy area of Talasari Beach (21°36'11"N, 87°24'27"E), Balasore, Odisha. This paper also reports the species for the first time from eastern coast of Indian Mainland as well as from Odisha state with its diagnostic characters. In India there are total thirteen species of genus *Anthopleura* have been reported without proper taxonomic study.

\*Corresponding author: Email: trideb400@gmail.com, trideb400sana@gmail.com;

Thus, this paper also deals with a comparative analysis of various sea anemone species of genus *Anthopleura* reported from Indian water.

**Keywords:** First record; Sea anemone; Actiniaria; *Anthopleura*; Indian mainland.

## 1. INTRODUCTION

Ocean comprises over 90 percent of the habitable space on the Earth and contains some 250,000 known species, with many more remaining to be discovered. At present, there are about 238,165 accepted marine faunal species described in the World Register of Marine Species database [1], yet this is only a fraction of all species living in the ocean, with new discoveries being made every year. India is one of the major mega-biodiverse countries which is home to a variety of species-rich ecosystems with an enormous 8,118 km, spread across the mainland and the islands of Andaman and Nicobar and Lakshadweep [2]. The marine ecosystem is home to a huge amount of marine faunal groups, although the sea anemone, order Actiniaria, Hertwig, 1882, are possibly one of the most understudied marine faunal groups. They are one of the notable benthic inhabitants whose occurrence have been reported across the all depths and latitudes [3,4,5]. Actinarians, popularly called as 'SeaAnemones', belongs to the phylum Cnidaria form a vital group of intertidal invertebrate differentiated by their habit, habitat and excellent colouration. The actinarian, sea anemones are gathered under the class Anthozoa that contains about 1107 described species occurring across the world [6]. This benthic faunal group is biologically very much vital for their close associations with various macrobenthic fauna such as clown fishes, shrimps and hermit crabs [7,8]. In a generalized view, sea anemones are cylindrical in shape; having three distinguished body part oral disc, column and pedal disc. The lower part of body bears pedal disc which provide site for firm attachment with solid substratum especially rock, wooden structure and dead coral. The upper end having mouth and the hollow tentacles arising from the oral disc surround it. The tentacle number varies species to species and arranged in a specific number of cycle which is the distinguished characters for species identification. The cylindrical column of the sea anemones is tapered towards the pedal disc bearing end and the oral disc is generally

greater in diameter. Sea anemones of India have had scant attention in comparison with other members of Anthozoa (especially scleractinian corals and octocorals).

The initiating works about sea anemones in Indian water were mainly focused on the occurrence and carried out by [9,10,11,12,13,14,15,16,17]. Later further study was conducted by multiple authors and everyone contributed to the science at their individual level. Revised checklist on the diversity of sea anemone of Indian water, prepared by Anushma et.al. [18] reports a total of 92 species of Actinarians under 22 families and 54 genera from Indian water. With special reference to genus *Anthopleura* there are total 13 species are reported so far from Indian water.

Present author came across a single specimen of genus *Anthopleura*, collected from Talasari Beach of Odisha; which is identified as *Anthopleura handi* Dunn, 1978. Previously the species was reported from Andaman Island, Bay of Bengal by Raghunathan et.al. [19]. So far there is no any report of this species from coast across the Indian mainland; therefore, this paper reports the occurrence of *Anthopleura handi* Dunn, 1978 for the first time from coast across Indian mainland. Previously there was a scarcity of literature regarding details taxonomic description of various species of genus *anthopleura* in Indian water. Here for the first time, we have added a comparative analysis of various sea anemone species of genus *anthopleura* reported from Indian water.

## 2. MATERIALS AND MATHODS

The specimen (n=1) was encountered while surveying across Talasari Beach (21°36'11"N, 87°24'27"E), Balasore, Odisha. It was observed that the column region of specimen was embedded inside the intertidal sandy area with tentacle exposed outside. After collection specimen it was preserved in 70% alcohol and brought to the laboratory for further stages of taxonomic work. Photography and the detailed



**Fig. 1. (A) Sample collection site with latitude and longitude; (B) Distribution of *Anthopleura handi* Dunn, 1978 throughout India**

morphometric analysis of the specimen was conducted. Photographs of specimens were taken in both live and preserved condition. Photograph of distinguishable characters of preserved specimens were taken under stereomicroscope, LEICA EZ4HD. Specimen was identified up to species level following the relevant literature [20,21]. Preserved specimens are deposited in the National Zoological Collections of General Non- chordate Section, Zoological Survey of India, Kolkata, India.

### 3. RESULTS

Taxonomic account of the species is given hereunder based on the collected specimen.

Phylum: **Cnidaria** Hatschek, 1888  
 Class: **Anthozoa** Ehrenberg, 1834  
 Order: **Actiniaria** Hertwig, 1882  
 Family: **Actinidae** Rafinesque, 1815  
 Genus: **Anthopleura** Duchassaing de fonbressin and Michelotti, 1860

**Material examined:** One example (Total length 19.28 mm, Total diameter 14 mm), Reg. No. ZSI/GNC/P4781/1, Collected by Santanu Mitra, collected on 02.11.2023 from Talsari Beach, Balasore, Odisha, India.

### 4. DIAGNOISTIC CHARACTERS

Specimen distinctly longer than broad (Length/diameter = 1.37) in preserved condition

(Fig.3). Column brownish yellow with gravels attached, 14 mm in diameter; oral disc grey; tentacles to 2-4 mm long in preserved specimen, with light bars across them (Fig. 2D); column with closely packed adhesive verrucae of 0.62-0.78 mm arrayed in a longitudinal manner (Fig. 2B); gravels are attached with verrucae (Fig. 2B); acrorhagus located at the distal end of each column of verrucae. Oral disc bearing lipless slit like mouth, slightly smaller than the pedal disc (Figs. 3 A, B). Tentacles equal in appearance, greyish yellow in colour in preserved condition, tapered bluntly at tip, approximately 60 in number, arranged irregularly (Fig. 2A), (Fig. 3 A, B); Mesenterial insertions visible through pedal disc as dark lines (Fig. 2C). Pedal disc roughly circular, adhesive and well organized (Fig.4).

**Habitat:** Specimens were attached to rocky substances or oyster shells on intertidal sandy flats, including the high inter-tidal regions, where the anemones inhabit sand around the base of rocks. It was found that when the base of specimen is attached to solid object or buried in sediment and oral disc was at or above surface of substratum.

**Colour in life:** Tentacles transparent with dark brownish horizontal stripes interrupted by narrow white cross bar at irregular interval throughout its entire length; column overall grey- brown with brownish - purple verrucae. Oral disc light greyish in colouration (Figs. 3 A, B). Acrorhagi

showing same colouration as column but some with white- tipped.

**Distribution (Within India):** Talsari, Odisha (Present record), Andaman Island, Bay of Bengal [19].

**Distribution (Outside India):** Malaysia; Singapore; Philippines; northern and eastern

Australia. Type locality: Jeram Beach, Strait of Malacca, Selangor, Penin - sular Malaysia.

**Remarks:** Previously reported from Andaman Island [14] but the occurrence of *Anthopleura handi* Dunn, 1978 from Odisha is a new distributional record to Indian mainland.




Fig. 2. (A) Oral end showing arrangement of tentacles around mouth, (B) Distribution of verrucae across the column, (C) Portion of pedal disc, (D) Tentacles



Fig. 3 (A), (B). Overall view of *Anthopleura handi* Dunn, 1978, (ZSI RegNo. ZSI /GNC/P4781/1) in living condition.




Fig. 4. *Anthopleura handi* Dunn, 1978 in preserved condition

## 5. DISCUSSION

Genus *Anthopleura* Duchassaing de Fonbressin & Michelotti, 1860 is represented by 102 species worldwide. Among them only thirteen species have been reported from Indian water. *Anthopleura handi* Dunn, [20] was described on the basis of 200 specimen originally. The present specimen collected by Santanu Mitra from Talsari Beach, Balasore, Odisha is almost similar

with the originally described specimen by Dunn, [20]. Original description mentioning the column may be drab grey, grey-brown, grey-green or tan in color but in case of present specimen column is grey-brown in colour. Original description also stating tentacle number ranges from ranges 45 to 61 whereas present specimen accounts for approximately 60 tentacles. A comparative analysis of all the species of genus *Anthopleura* reported from Indian water is given below;

List 1. Comparative analysis of all the species of genus *Anthopleura* reported from Indian water

Sl No.	Species Name	Tentacle	Distinguished Character (needed if any)
1	<i>Anthopleura annea</i> (Carlgren, 1940)	160 tentacles [22]	No
2	<i>Anthopleura asiatica</i> [23]	48 tentacles with greyish brown colouration [23]	oral disc greyish brown in colouration [23]
3	<i>Anthopleura buddiemieri</i> (Fautin 2005 )	100 tentacles Bijukumar et al. [24]	No
4	<i>Anthopleura dixoniana</i> (Haddon and Shackleton 1893)	60 tentacles [25]	Verruace greenish, brownish or orangish in colouration; Oral disc brown with green and yellow spot around the mouth [25].
5	<i>Anthopleura elegantissima</i> (Brandt 1835)	24 tentacles [26]	No
6	<i>Anthopleura handi</i> Dunn [20]	60 tentacles [20]	Oral disc greyish in colouration, column with closely packed adhesive verrucae of 0.62-0.78mm arrayed in a longitudinal manner throughout whole length [20]
7	<i>Anthopleura nigrescens</i> [27]	48-90 tentacles [27]	Oral disc dark gray-brown to black in colouration [27]
8	<i>Anthopleura panikkarii</i> [28]	96 tentacles [28]	No
9	<i>Anthopleura sola</i> (Pearse and Francis 2000)	Tentacle brownish with yellow band [29]	Oral disc green pigmentation [29].
10	<i>Anthopleura thalia</i> (Gosse, 1854)	30-60 tentacle; Tentacle greyish green to brownish yellow in colouration, slightly darker distally than proximally [30].	Verrucae are more prominent distally than proximally; Some specimens with red pigment overlying oral disc; red colouration especially concentrated near bases of tentacles and over the site of mesenterial insertions [30].
11	<i>Anthopleura waridi</i> (Carlgren, 1900)	Tentacles dark olive green to brown, tapered; arranged in up to five cycles [31].	No



Interestingly *Anthopleura handi* Dunn, 1978 can be distinguished from *A. annea*, *A. asiatica*, *A. buddiemieri*, *A. elegantissima*, *A. panikkarii*, *A. sola*, *A. thalia* and *A. waridi* on the basis of either number of tentacle or colour of tentacle but show some sort of similarity with *A. dixoniana* and *A. nigrescens* on the basis of tentacle number. In *A. handi* oral disc grey in colour whereas in *A. dixoniana* oral disc brown with green and yellow spot around the mouth [25] and in *A. nigrescens* oral disc dark gray-brown to black in colouration [27].

Interestingly the specimen was collected from Odisha, eastern coast of India and previously recorded from Andaman Islands [19]. Thus this finding has a great significance in geographical point of view as it is recorded for the first time from Indian mainland.

## 6. CONCLUSION

India shares a coastline of nearly 8,118 km that is exceptionally varied and dynamic and shows diverse rock-based, sediment-based and coral-dependent coastland landforms [2]. In worldwide level there are about 1107 described sea anemone species have been recorded [6] but in Indian water there are only 92 [18]. Data showing there is a huge lack of study on sea anemone in India. Sea anemone has a very unique importance in marine biodiversity as it acts as 'bioindicator'. Sea anemone shows mutualistic association with hermit crabs, mollusc, polychaetes and barnacle [32,33]. Thus on conservation purposes it has a significant value as it is associated with such a wide variety of macrofaunal invertebrate group. Previously the current species was reported from island biogeographic region but now it has been recorded from coastal biogeographic region and thus this finding shows a great value on geographical basis.

## ACKNOWLEDGEMENTS

Authors are thankful to Dr. Dhriti Banerjee, Director, Zoological survey of India, Kolkata for providing facilities and encouragement. We acknowledge Dr. L. K. Singh, Officer-in-Charge, F.P.S Building and Dr. N. Marimuthu, Officer-in-Charge, General Non-Chordata section for their technical help and encouragement during the present study.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. WoRMS Editorial Board World Register of Marine Species; 2024.  
Available: <http://www.marinespecies.org> at VLIZ Accessed January, 2024.
2. Gopalakrishnan A, Divya PR, Basheer VS, Raja Swaminathan T, Kathirvelpandian A, Bineesh KK, Kumar RG, Jena JK. Macro flora and fauna of the Gulf of Mannar - a checklist. National Bureau of Fish Genetics Resources. Lucknow, UP, India. 2012;127.
3. Ammons AW, Daly M. Distribution, habitat use and ecology of deep water Anemones (*Actiniaria*) in the Gulf of Mexico. Deep Sea Research Part II, Topical Studies in Oceanography, 2008;55(24):2657-266.  
DOI:10.1016/j.dsr2.2008.07.015.
4. Schories D, Reise K, Sanamyan K, Sanamyan N, Clasing E, Reise A. Actinian dominated intertidal mudflats: A new case of an extraordinary rare phenomenon from Southern Chile. Journal of Sea Research. 2011;65:304-314.
5. González-Muñoz R, Simões N, Guerra-Castro EJ, Hernández-Ortiz C, Carrasquel G, Mendez E, Lira C, Rada M, Hernández I, Pauls SM, Croquer A, Cruz-Motta JJ. Sea anemones (Cnidaria: Actiniaria, Corallimorpharia, Ceriantharia, Zoanthidea) from marine shallow-water environments in Venezuela: New records and an updated inventory. Marine Biodiversity Records. 2016;9:18.  
DOI: 10.1186/s41200-016-0016-7.
6. Fautin Daphne, Wallace Carden, Crowther Andrea. Fautin DG, Wallace CC, Crowther AL. Revision of the sea anemone fauna of Moreton Bay, S.E. Queensland. Memoirs of the Queensland Museum. Memoirs of the Queensland Museum 2008;54:35-64.
7. Bach CE, Herrnkind WF. Effects of predation pressure on the mutualistic interaction between the Hermit Crab, *Pagurus Pollicaris* Say, 1817, and the Sea Anemone, *Calliactis Tricolor* (Lesueur, 1817). Crustaceana. 1980;38(1)104-108.  
Available: <https://doi.org/10.1163/156854080X00508>
8. Fautin DG, Allen GR. Field guide to anemonefishes and their host sea

- anemones. Western Australian Museum, Francis Street, Perth; 1992.
9. Annandale N. The fauna of brackish ponds at Port Canning, Lower Bengal. P.1. Records of Indian Museum. 1907;1(4):45-74.
10. Stoliczka F. On the anatomy of *Sagartia schilleriana* and *Membranipora bengalensis*, a new coral and a bryozoan living in brackish water at Port Canning. Journal of Asiatic Society of Bengal. 1869; 38:28–63.
11. Annandale N. Fauna of Chilka Lake. The Coelenterates etc. Memoirs of Indian Museum. 1915;5:65-114.
12. Carlgren O. A revision of the *Actiniaria* of the Chilka Lake. Arkiv för Zoology. 1925; 17A:1-21.
13. Carlgren O. A survey of the Ptychodactaria Corallimorpharia and Actiniaria. *Kungl. Svenska vetenskapsakademiens handlingar*. 1949;1:1-121.
14. Menon KR. Subclass Zoantharia (except Scleractinia). Bulletin of Madras Govt. Museum (Natural History Section). 1927; 1(1):31-40.
15. Panikkar NK. The structure, bionomics and systematic position of two new brackish water Actiniaria from Madras. Proceedings of the Zoological Society of London. 1936; 106:39-52.
16. Panikkar NK, Aiyar RG. The brackish water fauna of Madras. Proceedings Section B Indian Academy of Science, Bangalore. 1937;6 (5):284-337.
17. Panikkar NK. Studies on *Peachia* from Madras. Proceedings Indian Academy of Science, Bangalore. 1938;7(4):182–205.
18. Anushma K, Prabhakaran MP, Deepa KP. A revised checklist of Sea *Anemones* (Order: Actiniaria) from the Exclusive Economic Zone (EEZ) of India. Journal of Aquatic Biology & Fisheries. 2022;10(1): 15-26.
19. Raghunathan C, Raghuraman R, Choudhury Smitanjali, Venkataraman Krishnamoorthy. Diversity and distribution of sea anemones in india with special reference to Andaman and Nicobar Islands. Records of the Zoological Survey of India. 2014;114:269-294.
20. Dunn DF. *Anthopleura handi* n. sp. (Coelenterata, Actiniaria), an internally brooding, intertidal sea anemone from Malaysia. Wasmann Journal of Biology. 1978;35:54-64.
21. Fautin DG, Wallace CC, Crowther AL. Revision of the sea anemone fauna of Moreton Bay, S.E. Queensland. Memoirs of the Queensland Museum. Memoirs of the Queensland Museum. 2008;54:35-64.
22. Den Hartog JC, Vennam J.. Some Actiniaria (Cnidaria: Anthozoa) from the West Coast of India. Zoologische Mededelingen, Leiden. 1993;67:601–637
23. Uchida T, Muramatsu S. Notes on Some Japanese Sea-anemones (With 5 Text-figures). Minutes of the Faculty of Science, Hokkaido University. 1958;14(1):111-119.
24. Kumar Biju. kv Dhaneesh, Geethu M. First record of the sea anemone *Anthopleura buddemeieri* fautin (Cnidaria: Actiniaria: Actiniidae) from the Indian coast. Journal of Aquatic Biology & Fisheries. 2015;3: 111-114.
25. Fautin DG, Tan SH, Tan R. Sea anemones (Cnidaria: Actiniaria) of Singapore: abundant and well-known shallow-water species. Raffles Bulletin of Zoology. 2009; 121-143.
26. Fautin DG, Hand C. Anthozoa. In: The light and smith manual: Intertidal invertebrates from central California to Oregon. J. T. Carlton (ed.). University of California Press, Berkeley. 2007;173-184.
27. Acuña, Fabián, González-Muñoz, Ricardo, Garese, Agustín, Díaz-Ferguson, Edgardo. First records of *Anthopleura nigrescens* (Verrill, 1928) and *Telmatactis panamensis* (Verrill, 1869) (Cnidaria, Anthozoa, Actiniaria) from Parque Nacional Coiba, Pacific coast of Panama. Check List. 2022; 18:1141-1146. DOI:10.15560/18.5.1141.
28. Parulekar AH. On a new species of sea anemone from Maharashtra, India. Journal of the Bombay Natural History Society. 1968;65(3):590-595,590-595.
29. Govenar K. Evidence of Photoenhancement by Green Fluorescent Proteins in Low Light Conditions: A Potential Factor of Habitat Distribution for the Sea Anemone *Anthopleura sola*. UC San Diego; 2021. ProQuest ID: Govenar\_ucsd\_0033M\_20222. Merritt ID: ark:/13030/m56j0m8s. Retrieved from: <https://escholarship.org/uc/item/0ws0d0fs>.
30. Daly Marymegan, Picton Bernard. Description of the Sea Anemone *Anthopleura Thallia* (Gosse 1854). Biology

- & Environment Proceedings of the Royal Irish Academy. 2012;112B:1-6.  
DOI:10.3318/BIOE.2012.10.
31. Gul, Shahnawaz. First records of sea anemones (Cnidaria: Actiniaria) from the coast of Pakistan (Northern Arabian Sea). Int. J. Biol. Biotech. 2017;14:247-252.
  32. Balakrishnan, Srinivasan, Santhanam, Perumal, Tudu Prasad, Yogesh Kumar JS, Mohapatra Anil. Mutualistic interactions of polychaete, barnacles, mollusca and sea anemone with hermit crab. Indian Journal of Geo-Marine Sciences. 2019;48:286-287.
  33. Antoniadou C, Vafeiadou AM, Chintiroglou C. Symbiosis of sea anemones and hermit crabs in temperate seas. Symbiosis: Evolution, Biology and Ecological Effects. NOVA Science, New York. 2013;95118.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://prh.mbimph.com/review-history/3173>