



Morphometric Analysis and Condition Factor of Freshwater Prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda) from the Gomti River, Lucknow, Uttar Pradesh, India

Samiksha Lodhi ^a, Aman Ahmed ^a, Veena P. Swami ^a
and Sanjive Shukla ^{a*}

^a P.G. Department of Zoology, BSNV PG College, University of Lucknow, Lucknow-226001 (U.P.), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.56557/UPJOZ/2024/v45i114071

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://prh.mbmph.com/review-history/3467>

Original Research Article

Received: 24/02/2024

Accepted: 29/04/2024

Published: 06/05/2024

ABSTRACT

The present investigations on morphometric analysis and condition factor of freshwater prawn, *Macrobrachium lamarrei* (in both sexes) from The Gomti river, Lucknow, Uttar Pradesh, India. The freshwater prawn, locally known as "Jhinga" was the subject of this study. Prawn samples had been collected in different months of both sexes, and a close relation between their length and weight were observed. The morphometric analysis showed that females ranged from 31-57 mm in length, with their weight ranges from 0.31-1.39g, which are larger than the males (21- 36mm) with their higher weight range (0.13- 0.49g). A high correlation coefficient was observed in males ($R^2 = 0.808$) and females ($R^2 = 0.881$) separately. There were significant differences in the condition

*Corresponding author: Email: sammylodhi841@gmail.com;

factors of males and females at $P = .05$. The lowest condition factor (CF) value for the males was observed in January (CF= 1.06) and highest in September (CF= 1.52), while in the females CF was the highest value in January (CF=0.82) and the lowest one in March (CF=0.75). The sex ratio of male: female (1.0: 2.10) was found, indicating that females are significantly greater in number than males.

Keywords: Freshwater prawn; *Macrobrachium* sp.; length-weight relationship; morphometric analysis; condition factor; the Gomti River.

1. INTRODUCTION

Crustaceans are an important component of the aquatic ecosystem. The shrimps, prawns, lobsters and crabs are a good source of protein for humans [1]. Crustacean Fisheries is an important growing industry of the world [2]. Nowadays more and more emphasis has been given to freshwater prawn culture across the globe [3,4]. Similar freshwater prawns like *Macrobrachium lamarrei* and *Macrobrachium dayanum* an “untapped resource”, are promising candidate for aquaculture having various advantages over marine prawn species as well as “Scampy” *Macrobrachium rosenbergii* [5].

The examination of the length- weight relationship plays a crucial role in the field of prawn biology. This investigation holds practical significance as it enables the evaluation of prawn growth in various environmental conditions [6]. The work on various crustaceans including freshwater prawn has also highlighted the importance of length-weight relationship analysis and condition factors to establish correlations among populations from different areas [7,8]. Furthermore, clarifying the morphometric analysis in aquatic animals, specifically fish, holds significant importance in fisheries and aquaculture [9]. It aids to determining the mean weight corresponding to a given length and also provides variations in growth rate of a specific species in different environmental conditions [10].

The CF (Condition Factor) of fish reveals the interactions between biotic and abiotic factors of the fish which reflects overall growth and development of animals [11]. Abowei [12] reported that the condition factor also reflects the influence of seasonal variations and habitat changes on the species robustness. Aquaculturists have documented the morphometric analysis and condition factor for prawns and fishes as crucial indicators of their growth, development and health [13]. Variability in these key parameters (morphometric analysis and condition factors) for prawns provides insights

into the suitability of aquatic ecosystems for prawn strength, standing stock, and their production [4,14].

Prawns and shrimps belonging to the genera *Macrobrachium* and *Panaeus* hold significant value worldwide, and are often used to enhance the flavour of food preparations due to their high protein content [15,16,17]. The freshwater prawn is a relatively small size compared to prawn species, readily available whole year in local habitats. It matures entirely in freshwater and holds promise for freshwater aquaculture [18,19,20,21]. Additionally, this prawn species serves as a bio-indicator for monitoring environmental pollution [22,23,24,25,26,27,28, 18,29,19].

Therefore, the present investigation focused on both morphometric and condition factor analysis of *M. lamarrei* from the river Gomti, Lucknow (U.P.), India.

2. MATERIALS AND METHODS

Freshwater prawns, *Macrobrachium lamarrei* (Fig. 2) were collected from different locations of the river Gomti, Lucknow (U.P.) (Fig. 1), with the help of local fisherman and brought to the laboratory (N- 26°49' 54" E- 80°55'58") and maintained in glass aquaria [30,22].

The total length of each individual was measured with the help of a divider, meter scale and vernier calliper, and the weight by digital weighing machine [31,32,33,34,35]. The male and female prawns were analysed separately by collecting them according to their gender, and by the month for body mass and length-weight relationship [36]. The length-weight relationship was calculated by linear regression [37] according to the following formula [38].

$$W = aL^b$$

Where:

W= Weight of Prawn(g)

L= Length of Prawn (cm)

a = Intercept; b= Regression coefficient




Fig. 1. Collection Site the Gomti river Lucknow (U.P) India (N-26°52'27" E- 80°54'42")



Fig. 2. Freshwater prawn, *M. lamarrei*

The parameters a and b were evaluated by linear regression [38,39,40,41,9]. This equation is sometimes also referred to as the length-weight key [42].

2.1 Condition Factor (CF)

It shows interaction between exogenous and endogenous factors which reflects the overall

growth and development of small prawns. It was calculated using the following formula [43]:

$$CF=100*W/L^3$$

Where:

CF= Condition Factor

W is the weight (g) of the prawn and L is the Total length of the prawn (cm)

2.2 Sex Ratio

By inspecting the presence or absence of appendix masculine in the second abdominal pleopod, the sex of the specimens was ascertained. The ratio of male and female was based on the frequency of the prawns caught during the sampling period. The sex ratio of the male and female individuals was calculated [44].

$$\text{Sex Ratio} = M/F$$

Where:

M = Number of male prawns
F = Number of female prawns

2.3 Statistical Analysis

The Chi square test was employed to assess the significant difference of sexes every month. All statistical analysis was done using Microsoft Excel 2019 and MINITAB Software.

3. RESULTS AND DISCUSSION

The Prawn samples were collected in different months for the both sexes to explain the relationship between their morphometric data. A total of 100 prawns had been taken as a sample size. The females were found throughout the year, but the males were not found in the February and March stocks. The males were found in lower numbers compared to females.

A strong positive significant relationship was observed between length and weight of *Macrobrachium lamarrei* ($R^2 = 0.881$) in July, and insignificant relationship was observed in the month of October ($R^2 = 0.356$). In the males, the relationship was significant in October ($R^2 = 0.808$) and weak relationship was observed in the month of September ($R^2 = 0.147$). Table 1 showed morphometric analysis between body weight and body length of female freshwater prawn, *Macrobrachium lamarrei* in different months. While Table 2 showed the morphometric relationship of the male prawns. The Sex ratio of the prawns showed significant difference in various months at ($P = .05$). Therefore, the result showed that the sexual frequency and sex ratio varied in different months. The highest sex ratio was observed in the month of September and the lowest one in February and March (Table 3).

The condition factor of the males showed the maximum value in September (CF = 1.52) and the minimum in January (CF = 1.06, Table 2)

whereas, in the females, the condition factor was highest in the January (CF = 0.83) and lowest in the March (CF = 0.75) (Table 1). A positive allometric growth of *Macrobrachium lamarrei* from the Gomti river on the basis of growth coefficient (b value) by using t-test ($P = .05$) (Tables 1 and 2). The values of growth coefficient were high in September, which indicates the suitability of the environment for food availability in the Gomti river [45]. Other species of *Macrobrachium* have also been reported to exhibit similar range of condition [46,43].

The study of the length-weight relationship of *M. lamarrei* showed a highly significant correlation between total length and body weight. The present study showed, the regression coefficient (b) for females is moderately high as compared to the males. This suggests that the females gain more weight with increase in the length, indicating better well-being [47], also reported the same observations in freshwater Prawn *Macrobrachium rosenbergii* in Earthen Pod at High Density (Ming Y et al., 2016 [48]. The regression coefficient values showed a positive allometric growth pattern, which corresponds to the outcomes of [49,50] regarding the *M. macrobrachion*, and [34] reported in *P. notialis*. [51], suggests that prawns become relatively robust and stronger as they increase in length when showing positive allometric growth patterns. Strong regression coefficient between length and weight were observed, with coefficients for males ($R^2 = 0.14-0.80$) and for females ($R^2 = 0.35-0.88$), indicating that the model is suitable for the study [32].

According to many factors such as food availability, feeding rate and reproductive activity the length-weight relationship may temporarily vary, however, the parameter 'b' is usually insignificant throughout the year [52]. In *M. dux* and *M. macrobrachion* the regression coefficient showed gender diversity for male and female prawns as reported by Iloba et al. [33].

Jimoh et al. [53] reported length- weight relationship of *Macrobrachium vollenhoveni* with regression coefficient value ranged from 2.58 to 3.11 from Badagry creek. The analysis of length-weight relationship was given by Le [38] in *Perca fluviatilis* [54,55], inspected the value of 'b' varying from 2.5 to 4.0. The evaluated parameter 'b' revealed negative allometric growth and was supported by high correlation values, suggesting elevated predictability between carapace length and weight of the studied shrimps [56, 57].

Table 1. Regression analysis and condition factor of the Female freshwater prawn, *Macrobrachium lamarrei*

Month	Weight (g) \pm SE	Length (mm) \pm SE	R ²	b	CF Value
January	0.83 \pm 0.09	45.7 \pm 0.13	0.768	1.25	0.83
February	0.81 \pm 0.05	46.9 \pm 0.10	0.528	1.37	0.78
March	0.58 \pm 0.03	42.5 \pm 0.08	0.635	2.09	0.75
April	0.83 \pm 0.08	46.2 \pm 0.16	0.815	1.76	0.81
May	0.69 \pm 0.06	44.7 \pm 0.16	0.761	2.14	0.76
June	0.89 \pm 0.07	47.7 \pm 0.11	0.366	0.87	0.81
July	0.76 \pm 0.05	45.2 \pm 0.12	0.881	2.06	0.81
August	0.83 \pm 0.08	46.7 \pm 0.14	0.660	1.37	0.79
September	0.88 \pm 0.07	46.9 \pm 0.11	0.812	1.39	0.82
October	0.73 \pm 0.05	45.8 \pm 0.12	0.356	1.34	0.77
November	0.87 \pm 0.06	47.1 \pm 0.11	0.806	1.53	0.81
December	0.77 \pm 0.06	45.5 \pm 0.11	0.803	1.63	0.80

Table 2. Regression analysis and condition factor of the Male freshwater prawn, *Macrobrachium lamarrei*

Month	Weight (g) \pm SE	Length (mm) \pm SE	R ²	b	CF Value
January	0.25 \pm 0.01	28.7 \pm 0.07	0.730	4.49	1.06
February	-	-	-	-	-
March	-	-	-	-	-
April	0.27 \pm 0.02	27.8 \pm 0.06	0.500	1.93	1.27
May	0.31 \pm 0.02	28.4 \pm 0.06	0.549	2.11	1.37
June	0.27 \pm 0.07	27.9 \pm 0.04	0.765	5.10	1.25
July	0.27 \pm 0.07	27.8 \pm 0.05	0.695	5.61	1.29
August	0.23 \pm 0.01	26.7 \pm 0.08	0.709	4.55	1.23
September	0.30 \pm 0.01	27.5 \pm 0.07	0.147	1.41	1.52
October	0.25 \pm 0.00	25.9 \pm 0.05	0.808	6.52	1.45
November	0.29 \pm 0.02	28.2 \pm 0.06	0.578	1.82	1.27
December	0.24 \pm 0.01	25.9 \pm 0.08	0.819	5.88	1.40

Table 3. Sex ratio of Male and Female freshwater prawn, *Macrobrachium lamarrei*

Month	M/F Ratio
January	0.28
February	0
March	0
April	0.36
May	0.25
June	0.31
July	0.28
August	0.42
September	0.47
October	0.38
November	0.28
December	0.31

In the present study, *Macrobrachium lamarrei* reflected the highest condition factor for the males (1.53 in September) and the females (0.83 in January). There were notable differences between male and female in different months (April, June, July, September) of the year. Similar results were observed by Aminisarteshnizi [32] in

M. nipponense in the Siah Darvishan River, [58] in *Metapenaeus monoceros*, Bhidiya and [50], in *M. macrobrachion* from the Kwa-river.

Kanaujia et al. [34] reported Fulton's condition factor for *Labeo rohita* ranged from 0.77506-2.45703. The key factors which affect the K value

consists of size, season, sex and development of gonads in fish [59]. The well-being of the fish is also influenced by food supply, ecological factors etc. [38,60,61,62] reported that the K value tends to decline with age. In Southern coasts of the Caspian Sea, Iran, condition factors of *M. nipponense* were found 0.89-1.90 [7,63].

The gender distribution in the present studies varies significantly from month to month. February and March showed a notable gender imbalance with 0 males and 100 females each. This suggests that these months might represent a specific stage in the prawn's life cycle or behavior, such as reproduction or molting when males are less active or not present.

4. CONCLUSION

Macrobrachium lamarrei shows a positive allometric growth in the Gomti river. The elevated value of regression coefficient "R" showed a strong relationship between total length and body weight. The high value of the coefficient of determination R^2 suggested that the model used for the analysis fits the data, hence confirming the fitness of the model. Knowledge of morphometric analysis and the condition factor of introduced or invaded species are essential to assess an appropriate management of alien and native species in an aquatic ecosystem. More advanced studies are required to determine where there is a correlation between each morphotype's physiological, behavioural and functional characteristics. An understandable explanation of the morphotype of this species is very important for understanding growth processes and adaptive value in the population to improve culture management.

ACKNOWLEDGEMENT

Authors are thankful to the Prof. U. D. Sharma, Rtd. Professor, Department of Zoology, University of Lucknow, Lucknow and also thankful to Head, P.G. Department of Zoology, B.S.N.V.P.G. College, Lucknow (U.P.) India, for providing necessary lab facilities.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Carnevali RP, Collins PA, Poi de Neiff ASG. Trophic ecology of the freshwater prawn, *Pseudopalaemon bouvieri* (Decapoda: Palaemonidae) in Northeastern Argentina, with remarks on population structure. *Rev. Biol. Trop.* 2012;60(1):305-316.
2. Bondad- Reantaso MG, Subasinghe RP, Josupeit. H, Cai J, Zhou X. the role of crustacean fisheries and aquaculture in global food security: Past, present and future. *Journal of Invertebrate Pathology.* 2021;110(2):158-165.
3. Mohanty KR, Sunderray KJ. Environmental impacts of coastal shrimp farming and measures to reduce consequential pollution. *Fishing Chimes.* 1999;19:7.
4. Sankar G, Elavarasi A, Sakkaravarthi K, Ramamoorthy K. Biochemical changes and growth performance of Black tiger Shrimp larvae after using Ricinus communis extracts as feed additive. *Intl. J Phar Tech. Res.* 2011;3;201-208.
5. Shukla S, Sharma UD. Smaller fresh water prawns: Their aquaculture potential and suitability as good laboratory model. In: Bioresources for food security and rural livelihood. Kulkarni GK, Pandey PN. (Eds). Narendra Publications, Delhi. 2010;189-204.
6. Bahuguna P, Joshi HK. Statistical observation on the length-weight relationship of brain and body in a coldwater catfish *Amblyceps mangois* (Ham. -Buch.) from Garhwal region. *Environ. Conser. Journal.* 2010;11(1-2): 21-23.
7. Mossolin EC, Bueno SLS. Relative growth of the second pereopod in *Macrobrachium olfersi* (Wiegmann, 1836) (Decapoda, Palaemonidae). *Crustaceana.* 2003;76(3): 363-376.
8. Pantaleão JAF, Hirose GL, Costa RC. Relative growth, morphological sexual maturity, and size of *Macrobrachium amazonicum* (Heller 1862) (Crustacea, Decapoda, Palaemonidae) in a population with an entirely freshwater life cycle. *Invertebrate Reproduction and Development.* 2012;56(3):180-190.
9. Khanipour AA, Noori A, Amini M, Kamrani, E. Length-weight relationship and Fulton's condition factor of *Macrobrachium nipponense* (De Haan, 1849) in Anzali lagoon of Iran. *Iranian Journal of Fisheries Sciences,* 2020;19(1):496-500.
10. Abowei JFN, Tawari C, Cdeekae SN, Amakiri NE. A study of the length –weight relationship and condition factor of

- Pseudotolithus elangatus 559 Fulton's Condition Factor of Macrobrachium nipponense in Siah Darvishan River (Browdich, 1825) from Bonny Estuary, Niger Delta, Nigeria. Int. J. Trop. Agr. Food.Sys, 2008;2(3-4):249-254.
11. Rajeevan R, Edirisinghe U, Athauda ARSB. Length- weight relationships and Condition factor of giant freshwater prawn, Macrobrachium rosenbergii (De Man, 1879) in five perennial reservoirs in Northern Province, Sri Lanka. International Journal of Fisheries and Aquatic Studies. 2018;6(5):283-287.
12. Abowei JFN. The condition factor, length - weight relationship and abundance of Ilisha africana (Block, 1795) from Nkoro River Niger Delta, Nigeria. Adv. J. Food Sci. Technol. 2010;2(1):6-11.
13. Attia AO. El-Aiatt, Kariman A. Sh. Shalloof. Study on the biology of the small scaled terapon Terapon puta (Cuvier, 1829) from Bardawil lagoon, North Sinai, Egypt. Egyptian Journal of Aquatic Biology and Fisheries. 2019;23(2):95 – 107.
14. Shallof KA. Sh, El-Far AM. Length-Weight Relationship and Condition Factor of Some Fishes from the River Nile in Egypt with Special Reference to Four Tilapia Species. Egyptian Journal of Aquatic Biology and Fisheries. 2017;21(2):33-46.
15. Adeosun FI, Adedoyin IA, Oghenebrohie OMT, Quddus AA, Micheal AO. Morphometric and Meristic features and Length-weight Relationship as indicators of quality of Brycinus macrolepidotus in Lower River Ogun, Nigeria. Egyptian Journal of Aquatic Biology and Fisheries. 2019;23(2):433-441.
16. Cruz YM, Olatunbosun O. Comparative Study on the efficiency of three different types of crab pot in Iceland fishing ground. UNU-Fisheries Training Program. 2013;4:1-25.
17. Major RN, Taylor DI, Conner S, Conner G. Factors affecting bycatch in a developing New Zealand scampi potting fishery. Fisheries Research. 2016;10:55-64.
18. Jayachandran KV, Indira. Prawn Fisheries Resources of India for food security as well as for rural employment. 97th Indian Sci. Congr. Symposium, January 3-7, Thiruvananthapuram; 2010.
19. Shukla S, Sharma UD. Smaller fresh water prawns: Their aquaculture potential and suitability as good laboratory model. In: Bioresources for food security and rural livelihood. Kulkarni GK, Pandey PN. (Eds), Narendra Publications, Delhi. 2010;189-204.
20. Shukla S, Shukla R, Shukla S, Ahmad A, Mishra A. Effect of temperature on chromatophores of freshwater prawn, Macrobrachium lamarrei (Crustacea-Decapoda). Anusandhaan. 2017;5(1):80-84.
21. Ahmed A, Lodhi S, Shukla S. Observations on feeding behaviour of fresh water prawn, Macrobrachium lamarrei (Crustacea: Decapoda) Int. J. of Fish. and Aqua. Stud. 2021;9(6):109-112.
22. Lodhi S, Ahmed A, Lodhi HS, Shukla SA report on freshwater prawn, Macrobrachium lamarrei (Crustacea: Decapoda) from river Gomti, Lucknow (U.P.), India. Journal of Experimental Zoology. India. 2024; 27(1):751-756.
23. Sharma UD, Shukla S. Behaviour dysfunction of fresh water prawn, Macrobrachium lamarrei (Crustacea-Decapoda) following exposure to synthetic detergent, linear alkyl Benzene Sulphonate. Biol. Mem. 1990;16(12):58-61.
24. Sharma UD, Shukla S. Acute toxicity of heavy metals and detergent to fresh water prawn, Macrobrachium lamarrei (Crustacea-Decapoda). Him. J. Env. Zool. 2006;20(1):1-6.
25. Lodhi HS, Khan MA, Verma RS, Sharma UD. Acute toxicity of Copper Sulphonate to freshwater prawns. J. Environ. Biol. 2006; 27(3):585-588.
26. Lodhi HS, Khan MA, Verma RS, Sharma UD. Seasonal fluctuations in the total haemocyte counts (THC's) of freshwater prawn, Him. J. Env. Zool. 2004;18(1): 1-5.
27. Lodhi HS, Tiwari KJ, Shukla S, Sharma UD. Copper induced fluctuations in total haemocyte counts (THCs) of freshwater prawn, Macrobrachium lamarrei (Crustacea-Decapoda). J. Env. Bio. Sci. 2008;22(2):135-142.
28. Lodhi HS, Shukla S, Sharma UD. Studies on structure of heart of two freshwater prawns under light microscope. Uttar Pradesh J. Zool. 2009;29(1):1-10.
29. Verma DR, Lodhi HS, Tiwari KJ, Shukla S, Sharma UD. Copper sulphate induced changes in scaphognathite oscillations and oxygen consumption of fresh water prawn, Macrobrachium lamarrei (Crustacea-Decapoda). Journal of Applied and Natural Science. 2010;2(1):34-7.

30. Sharma M, Lodhi S, Ahmed A. Freshwater prawn aquaculture: Prospects in U.P. Int. J. of Fish. and Aqua. Stud. 2022;10(4):146-151.
31. Lima JDF, Da Cruz MCM, Silva LMAD. Reproductive biology of *Macrobrachium surinamicum* (Decapoda: Palaemonidae) in the Amazon River mouth. Acta Amazonica. 2015;45(3):299 – 306.
32. Aminisarteshnizi. M. Length -weight relationship and Fulton's Condition Factor of *Macrobrachium nipponense* (De Haan, 1894) in Siah Darvishan River, Iran. Egyptian Journal of Aquatic Biology and Fisheries. 2021;25(2):551-560.
33. Iloba, Isioma. K, Otokuefor, Zino. J, Israel OS. Length-weight relationship and health of *Macrobrachium dux* (Lenz, 1910) and *Macrobrachium macrobrachion* (Herklots, 1851) in river Ethiopie Headwaters, Nigeria. Uttar Pradesh Journal of Zoology. 2022;43(5):1-13.
34. Kanaujia S, Sahu A, Verma DK, Singh MB. Length-weight relationship and condition factor of a commercial food fish *Labeo rohita* (Hamilton, 1822) from Ghaghara River, Uttar Pradesh, International Journal of Fisheries and Aquatic Studies. 2023;11(4):96-99.
35. Ibrahim S, Zhong Z, Lan X, Luo J, Tang Q, Xia Z, Yi S, Yang G. Morphological Diversity of Different Male Morphotypes of Giant Freshwater Prawn *Macrobrachium rosenbergii* (De Man, 1879). Aquaculture Journal. 2023;3:133-148.
36. Bahuguna P, Dimri A, Rayal R, Sharma N. Observations on the body mass weight-length relationship and relative condition factor of *Macrobrachium assamensis peninsularis* (Tiwari) from Khoh river, Uttarakhand, India. Uttar Pradesh Journal of Zoology. 2021;43(13):54-65.
37. Pauly D. Some simple methods for the assessment of tropical fish stocks. FAO fish tech Rap; 1983;234.52.
38. Le Cren ED. The length-weight relationship and seasonal cycle in gonad weight and condition in Perch (*Perca fluviatilis*). J. Anim. Ecol. 1951;20:201-219.
39. Jones RE, Petrell RJ, Pauly D. Using modified length-weight relationships to assess the condition of fish. Aquacultural Engineering, 1999;20:261-276.
40. Koutrakis ET, Tsikliras AC. Length- weight relationships of fishes from three northern Aegean estuarine systems (Greece). J. Appl. Ichthyol. 2003;19:258-260.
41. Sparre P, Ursin E, Venema SC. Introduction to tropical fish stock assessment. Part 1. Manual FAO Fisheries Technical. FAO Rome. 1989;337.
42. Biswas SP. Length-weight relationship and condition factor. In: Manual of methods in Fish Biology. South Asian Publishers, New Delhi. 1993;60-64.
43. Enin U. Length- weight parameters and condition factor of two West African Prawns. Rev. Hydrobiol. Trop. 1994;27:121-127.
44. Peña-Mendoza B, Gómez-Márquez JL, Salgado-Ugarte IH, Ramírez-Noguera D. Reproductive biology of *Oreochromis niloticus* (Perciformes: Cichlidae) at Emiliano Zapata dam, Morelos, Mexico. Rev. Biol. Trop, 2015;53(3-4):515-522.
45. Ara MG, Nobi MN, Fatima MK, Ahmed ZF. Evaluation of condition factor of a small indigenous freshwater prawn, *Macrobrachium lamarrei* (H. Milne Edwards, 1837) in Bangladesh. International Journal of Natural and Social Sciences. 2014; 1(2):71-76.
46. Arimoro. FO, Meye JA. Some aspects of the biology of *Macrobrachium dux* (lenz,1910) (crustacea: Decapoda: natantia) in river Orogondo, Niger delta, Nigeria. Acta Biologica Columbiana. 2007; 12(1):111-122.
47. Bahuguna P, Shah KK, Kumar, R. Observation on the length-weight relationship and relative condition factor of *Barilius Bendelisis* (Ham.) inhabiting a spring-fed tributary of river Alaknanda (Garhwal Himalaya), India. J.Natcon. 2009; 21(2):215-220.
48. Ming Y, Fujiang D, Xilin DAI. Length-length, Length-weight Relationships and Condition Factor of Giant Freshwater prawn, *Macrobrachium rosenbergii* (de Man 1879) Cultured in Earthen Pond at High Density. Journal of Asian Fisheries Science. 2016;29:164-177.
49. New MB, Valenti WC, Tidwell IH, D'Abram OL, Kuttay MN. Freshwater prawns: Biology and farming. Willy – Blackwell Publishing Ltd; 2010.
50. Andem AB, Idung JU, George E, Ubong GU. Length-weight relationship and Fulton's Condition Factor of brackish river prawn (*Macrobrachium macrobrachion*, Herklots, 1851) from Great Kwa River, Obufa Esuk beach, Cross River state, Nigeria. European Journal of Experimental Biology, 2013;3(3):722-730.

51. Lawal-Are AO, Akinjogunla VF. Length-Weight Relationships and Condition Factor in Lagos Lagoon, South West, Nigeria. *Science and Technology*. 2012;2(3):32-40.
52. Begenal TB, Tesch FW. Age and growth in methods of assessment of fish production in freshwater, Ed. Bagenal, T. Oxford. Blackwell Scientific Publication. 1978;101-136.
53. Jimoh AA, Fakoya KA, Hammed AM, Amosu AO, Kumolu- Johnson CA. Meristic and Morphometrics in the African River Prawn, *Macrobrachium vollenhovenii* (Herklots, 1857) from Ologe Lagoon, South West, Nigeria. *Journal of Agriculture and Environmental Research Studies*. 2005;1(1):12-18.
54. Hile R. Age and growth of the Cisco, *leucichthys artedti* (Le seum) in the North-Eastern high lands. *Wisconsin Bull U.S. bur. Fish*. 1936;48:211-317.
55. Martin WR. The mechanics of environmental control of body form in fishes. *Univ. Toronto Stud. Biol*. 58: Publ. Ant. Fish Res. Lab. 1949;70:1-91.
56. Carlander KD. Handbook of freshwater fishery biology. Ames. Fishing News Books. 1969;1.
57. Froese R. Cube Law. Condition factor and weight-length relationship: History, meta-analysis and recommendations. *J. Appl. Ichthyol*. 2006;22:241-253.
58. Dineshbabu AP. Length-weight relationship and growth of the speckled shrimp *Metapenaeus monoceros* (Fabricius) off Saurashtra. *Journal of Marine Biology Ass. India*, 2006;48(2): 180 - 184.
59. Heincke F. Bericht über die Untersuchungen der Biologischen Anstalt auf Helgoland zur Naturgeschichte der Nutzfische. (1. April 1905 bis 1. Oktober 1907). [Report on the investigations of the Biological Institute on Helgoland on the natural history of commercial fish (April 1, 1905 to October 1, 1907). In: Die Beteiligung Deutschlands an der Internationalen Meeresforschung, 4. & 5. Jahresbericht. Verlag von Otto Salle, Berlin; c1908.
60. Tenali DR, Phom NP, Yamala SP, Gudipudi SB, Kashyap S, Nirmalkar R. Evaluation of Length-weight relationship and condition factor among the cohorts of *Labeo calbasu* in Pachipenta reservoir, Andhra Pradesh. *Journal of Experimental Zoology*. 2023;26(2):2329-2332.
61. Gandotra R, Vivek Singh D, Shanker R. Length-weight relationship and condition factor of *Aspidoparia morar* (Ham.) inhabiting river Tawi and its tributaries. *Biosciences, Biotechnology Research Asia*. 2009;6(2):767-772.
62. Gandotra R, Shanker R, Ahmed S, Sagar S. Morphometry of *Tor putitora* (Ham.) from Jhajjar stream, Jammu (J&K). *J. Inland Fish Soc. India*. 2008;40(1): 86-89.
63. Namin JI, Nami E, Heidary S. Length-Weight Relationship and Fulton's Condition Factor of *Macrobrachium nipponense* (Dehaan, 1849) in southern coasts of the Caspian Sea-Iran. *International journal of Advanced Biological and Biomedical Research*. 2014;2(5):1650-1656.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://prh.mbimph.com/review-history/3467>