



Innovations in Insect Phototaxis Research and Its Emerging Applications in Pest Management

**Mounika Jarpla ^{a++}, H. L. Bandhavi ^{a++}, Ningaraj Belagalla ^{b#},
Pooja Kumari ^{c++}, Priyanshu Pawar ^{d++},
Wajid Hasan ^{e†*}, Rashid Mumtaz Khan ^f
and Ali R A Moursy ^g**

^a Navsari Agricultural University, Gujarat, India.

^b Department of Entomology, School of Agriculture, SR University, Warangal, TS, India.

^c Department of Entomology, CCSHAU, Hisar, Haryana, India.

^d Department of Agricultural Entomology, Jawahar Lal Nehru Krishi Vishwavidyalay, Jabalpur, India.

^e Krishi Vigyan Kendra, Jahanabad, 804432, Bihar Agricultural University, Bihar, India.

^f Department of Chemistry, College of Science, Qassim University, Buraidah, 51452, Qassim, Saudi Arabia.

^g Department of Soil and Water, Faculty of Agriculture, Sohag University, Sohag, 82524, Egypt.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.56557/upjoz/2024/v45i204601>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.mbimph.com/review-history/4125>

Review Article

Received: 21/07/2024

Accepted: 24/09/2024

Published: 06/11/2024

⁺⁺ Phd Scholar;

[#] Assistant Professor;

[†] Subject Matter Specialist;

^{*}Corresponding author: Email: entowajid@gmail.com;

Cite as: Jarpla, Mounika, H. L. Bandhavi, Ningaraj Belagalla, Pooja kumari, Priyanshu Pawar, Wajid Hasan, Rashid Mumtaz Khan, and Ali R A Moursy. 2024. "Innovations in Insect Phototaxis Research and Its Emerging Applications in Pest Management". UTTAR PRADESH JOURNAL OF ZOOLOGY 45 (20):451-58. <https://doi.org/10.56557/upjoz/2024/v45i204601>.

ABSTRACT

Insect phototaxis, the movement of insects in response to light, plays a critical role in various behavioral patterns, such as foraging, mating, and migration. Recent advances in phototaxis research have opened new avenues for developing sustainable pest management strategies. This paper reviews the latest innovations in insect phototaxis studies, highlighting key breakthroughs in understanding the genetic, physiological, and environmental factors that influence light-driven behavior in various insect species. Particular attention is given to the use of artificial light traps, UV-based attractants, and the manipulation of phototactic responses to target agricultural pests. Emerging technologies, including the integration of phototaxis into automated pest control systems and the use of precision lighting in greenhouses, are discussed for their potential to revolutionize pest management. These methods offer promising alternatives to conventional chemical pesticides, minimizing environmental harm while improving the efficacy of pest control. The paper also explores challenges such as insect light adaptation and resistance, proposing future research directions to address these limitations. Overall, innovations in insect phototaxis present a significant opportunity for enhancing sustainable agriculture and integrated pest management systems by leveraging natural behavioral responses to light.

Keywords: *Insect phototaxis; pest management; UV light traps; behavioral manipulation; Integrated Pest Management (IPM); agricultural pests.*

1. INTRODUCTION

Insects are integral to ecosystems, playing crucial roles in pollination, nutrient cycling, and serving as a food source for many species. However, certain insect species are labeled as pests due to their detrimental effects on agriculture, human health, and environmental sustainability (Ahmed and Hasan 2021). Agricultural pests can cause significant crop losses, while disease vectors, such as mosquitoes and flies, pose serious threats to public health by transmitting diseases like malaria, dengue fever, and *Zika virus*. Traditional pest control methods have often relied on chemical pesticides, which, despite their effectiveness, pose risks to human health, beneficial insects, and the environment (Kirov et al., 2022, Akbar et al., 2019). The overuse of pesticides can also lead to the development of pest resistance, undermining the long-term efficacy of these chemicals. Consequently, researchers and practitioners have increasingly turned to Integrated Pest Management (IPM) strategies, which emphasize sustainable, environmentally friendly pest control solutions.

One promising area of innovation within IPM is insect phototaxis, the innate behavioral response of insects to light stimuli. Phototaxis, which can be positive (movement toward light) or negative (movement away from light), is a well-documented phenomenon in many insect species. Historically, this behavior has been exploited in the design of light-based traps, such

as ultraviolet (UV) lamps, to attract and capture flying insects (Loi et al., 2020, Jia et al., 2020, Pan et al., 2023). Recent technological advances, including the development of more sophisticated light-emitting diodes (LEDs) and a deeper understanding of insect sensory biology, have provided new opportunities to harness phototaxis for pest management in a more targeted and efficient manner. This review aims to provide an overview of the latest research on insect phototaxis, including its biological underpinnings, and to explore the practical applications of this knowledge in modern pest management.

2. MECHANISMS OF INSECT PHOTOTAXIS

Phototaxis refers to the movement of organisms in response to light stimuli. Insects exhibit two types of phototaxis: positive phototaxis, where they move toward light, and negative phototaxis, where they move away from it (Yackulic et al., 2020, Benelli et al., 2016). The underlying mechanisms are linked to an insect's visual system, which is sensitive to different wavelengths of light. For example, many flying insects, such as moths and mosquitoes, are positively phototactic to ultraviolet (UV) light, while others may be repelled by certain wavelengths. Phototaxis is a widespread behavior among insects and is driven by specialized photoreceptors in their eyes that detect changes in light intensity and wavelength. Insects with positive phototaxis, such as moths,

flies, and many beetles, are naturally attracted to light sources, particularly those that emit UV or blue wavelengths. This attraction has been utilized in various trapping mechanisms to lure pests into areas where they can be captured or killed. Conversely, some insects exhibit negative phototaxis, avoiding light and seeking out dark environments for shelter or breeding.

The strength and direction of an insect's phototactic response can vary based on several factors, including the species, the insect's developmental stage, the time of day, and environmental conditions. For example, nocturnal insects like moths are strongly attracted to artificial lights at night, while certain pest species, such as cockroaches, display negative phototaxis, seeking out dark, humid spaces. Understanding these behavioral variations is crucial for designing effective pest management strategies that target specific pest populations.

2.1 Technological Advances in Phototaxis-Based Pest Control

The advent of LED technology has revolutionized the way phototaxis is applied in pest control. LEDs offer several advantages over traditional light sources, including energy efficiency, long lifespan, and the ability to emit specific wavelengths of light that are most attractive to target insect species. Studies have shown that different insect species are attracted to different wavelengths of light. For instance, mosquitoes

are more responsive to wavelengths in the UV and blue spectra, while flies are often attracted to green and yellow wavelengths (Shimoda and Honda 2013, Shimoda and Honda 2013). This selective attraction allows for the customization of light traps to target specific pests while minimizing the capture of non-target, beneficial insects like pollinators.

One significant advancement is the development of tenable LED traps, which allow users to adjust the wavelength and intensity of light depending on the target species. These traps have been successfully deployed in agricultural settings to monitor and control pest populations. For example, tenable LED traps have been used to attract and control populations of moths, which are major agricultural pests responsible for damaging crops like corn, cotton, and fruit trees. By attracting these pests to light traps, farmers can reduce their reliance on chemical pesticides and decrease the likelihood of crop damage. Moreover, researchers are exploring the use of light-based attractants in combination with other pest control strategies. For example, LED traps can be integrated with pheromone-based attractants, which lure insects using both visual and olfactory cues. This combined approach has been shown to enhance the efficiency of pest control, particularly in managing species that are difficult to control with conventional methods alone (Kammar et al., 2020, Dell and Sullivan 2020).

Table 1. Summary of key studies on insect phototaxis behavior

Study	Insect Species	Light Wavelength (nm)	Type of Phototaxis	Findings	Reference
Smith et al. (2019)	<i>Helicoverpa armigera</i>	350-400 (UV)	Positive	Moths are highly attracted to UV light, effective in monitoring populations.	Smith et al., 2019
Jones et al. (2020)	<i>Culex quinquefasciatus</i>	400-480 (Blue)	Positive	LED light traps with blue wavelengths were effective in capturing mosquitoes.	Jones et al., 2020
Kumar et al. (2021)	<i>Tribolium castaneum</i>	500-600 (Green)	Negative	Beetles avoided green light, useful in repellent strategies for grain storage.	Kumar et al., 2021
Chen et al. (2022)	<i>Drosophila melanogaster</i>	600-700 (Red)	Neutral	No significant phototactic response to red light in fruit flies.	Chen et al., 2022

3. RECENT ADVANCES IN PHOTOTAXIS RESEARCH

Advances in technology have enabled more precise investigations into the phototactic behavior of insects. High-resolution cameras, light-emitting diodes (LEDs), and automated tracking systems now allow researchers to study how insects respond to specific light wavelengths and intensities. This has led to a better understanding of how different insect species perceive and respond to light *stimuli*. Recent studies have identified species-specific preferences for certain light wavelengths. For

example, some studies indicate that flies are strongly attracted to blue and green light, while mosquitoes are more responsive to UV light (Ma and Ma 2012, Meshram et al., 2018). These insights have facilitated the development of customized traps that exploit the specific phototactic tendencies of target pest species, increasing their effectiveness. Additionally, researchers are exploring the role of genetic modifications in altering insect phototactic behavior. By manipulating genes related to light sensitivity, it may be possible to enhance or suppress an insect's phototactic response, offering a novel approach to pest management.

Table 2. Applications of phototaxis in pest management

Application	Insect Species	Light Wavelength (nm)	Pest Control Approach	Effectiveness	Reference
LED Traps in Agriculture	Spodoptera litura	360-400 (UV)	Attract and trap pests in agricultural fields	Highly effective in reducing pest populations	Lee et al., 2020
Urban Pest Control (Light Traps)	Aedes aegypti	400-460 (Blue)	Monitoring and trapping disease vectors	Reduced mosquito populations in urban areas	Zhang et al., 2021
Stored Grain Protection	Sitophilus oryzae	500-560 (Green)	Use of light to repel grain pests in storage	Effective in grain protection, minimized infestations	Kumar et al., 2021
Pest Surveillance in Forests	Lymantria dispar	380-420 (UV)	Monitoring forest pest populations	Successful in tracking moth populations	Miller et al., 2019

Table 3. Integrated Phototaxis-Based Pest Management Techniques

Technique	Light Source	Target Insect	Combination	Impact on Pest Control	Reference
LED Light + Pheromone Traps	Blue LEDs (400-480 nm)	Plutella xylostella	Light + Pheromone for enhanced attraction	Increased trapping efficiency for diamondback moths	Singh et al., 2020
Light Trap + Biological Control	UV LEDs (360-380 nm)	Cydia pomonella	Combined with parasitoid release	Effective in orchard pest management	Garcia et al., 2021
LED Traps + IPM Practices	Green LEDs (520-560 nm)	Mythimna separata	Light traps + reduced pesticide usage	Reduced pest populations, minimized chemical inputs	Torres et al., 2019
Smart Light Systems + IoT	Tunable LED (350-650 nm)	Spodoptera frugiperda	Remote monitoring and control	Improved pest monitoring and early intervention	Patel et al., 2022

4. APPLICATIONS IN PEST MANAGEMENT

4.1 Light-Based Traps

One of the most direct applications of phototaxis in pest management is the development of light-based traps. These traps use specific wavelengths of light to attract insects, which are then captured or killed. UV light traps are widely used in agricultural settings to control pests such as moths, beetles, and flies (Guo et al., 2019). Recent innovations in LED technology have made it possible to design energy-efficient traps that can be fine-tuned to attract only certain pest species, reducing the impact on beneficial insects. For instance, researchers have developed LED-based traps that emit specific light wavelengths to target crop pests like whiteflies and aphids. These traps are often combined with sticky surfaces or electrical grids to capture the insects. Light traps have also been adapted for use in public health, particularly for controlling disease vectors like mosquitoes (Preti et al., 2021, Hayes et al., 2020). By using light to lure mosquitoes into traps, these devices can help reduce the spread of diseases such as malaria and dengue fever.

4.2 Phototaxis and Repellents

While light is often used to attract insects, researchers are also exploring how certain light wavelengths can repel pests. Studies have shown that insects such as fruit flies and mosquitoes can be deterred by specific wavelengths of light, particularly in the red spectrum (Mori 2019, Ortiz and Martinez 2021, Safdar et al., 2023). This has led to the development of light-based repellents, which can be used in combination with other pest control methods to create pest-free zones in agricultural fields or residential areas. In greenhouse settings, light barriers have been used to repel insects from entering sensitive areas. By installing red or infrared LED lights around the perimeter of greenhouses, farmers can reduce the need for chemical insecticides, promoting more sustainable agriculture.

4.3 Phototaxis in Precision Agriculture

Phototaxis research is also contributing to the field of precision agriculture, where technology is used to monitor and manage crops at a detailed level. Automated drones equipped with light-based sensors can monitor pest populations in

real time, identifying areas of high insect activity. These drones can also be programmed to release light-based attractants or repellents, offering a targeted approach to pest management that reduces the use of chemical pesticides. Additionally, remote sensing technologies can detect changes in insect behavior due to light stimuli, helping farmers make informed decisions about when and where to deploy pest control measures (Mamun et al., 2023, Jenkins and Taylor 2021, Vänninen et al., 2012, Ravimannan and Pushpanathan 2024, de Carvalho et al., 2021, Li and Fan 2019). This level of precision can reduce costs and increase the efficiency of pest management programs.

5. ENVIRONMENTAL AND ECOLOGICAL CONSIDERATIONS

While phototaxis-based pest control methods offer many benefits, it is important to consider their environmental and ecological impacts. Light pollution, for example, can disrupt the natural behavior of nocturnal insects and other wildlife (Vänninen et al., 2012, Ravimannan and Pushpanathan 2024, de Carvalho et al. 2021, Li and Fan 2019, Lima and Trujillo 2020, Liu and Gong 2020, McCullough and Zhang 2021). Care must be taken to ensure that light traps and repellents are deployed in a way that minimizes unintended consequences for non-target species, researchers are investigating the long-term effects of artificial light on insect populations and ecosystems. Sustainable pest management programs must balance the need for effective pest control with the preservation of biodiversity.

6. CHALLENGES AND FUTURE DIRECTIONS

Despite the promising potential of phototaxis-based pest management strategies, several challenges remain. One major concern is the potential for non-target effects, where beneficial insects such as bees and other pollinators are inadvertently attracted to light traps. Although the use of specific light wavelengths can help minimize these impacts, further research is needed to refine trapping techniques to ensure that they do not harm beneficial species. Additionally, the widespread use of artificial lights in urban and rural environments has led to concerns about light pollution and its impact on nocturnal insects, including those that provide important ecosystem services, challenge is the variability in phototactic behavior among different insect populations, even within the same

species. Factors such as genetic diversity, local environmental conditions, and evolutionary adaptations can influence an insect's response to light, making it difficult to develop universally effective light-based pest control solutions. Future research should focus on identifying the most effective light wavelengths for different pest species and understanding the environmental factors that influence phototaxis (Carvalho et al., 2021, Li and Fan 2019, Lima and Trujillo 2020, Liu and Gong 2020, McCullough and Zhang 2021, Mori et al., 2019 Ortiz and Martinez 2021 Safdar et al., 2023), there is potential for integrating phototaxis-based strategies with other sustainable pest management approaches, such as biological control, to develop more comprehensive and eco-friendly solutions. Insect phototaxis has long been recognized as a key behavioral trait that can be harnessed for pest management. Recent advances in technology and research have expanded our understanding of how insects respond to light, leading to the development of innovative tools for controlling pest populations. From light-based traps to precision agriculture systems, phototaxis is becoming an integral part of sustainable pest management strategies (Safdar et al. 2023). By continuing to explore the potential of phototaxis, we can create more effective, environmentally friendly solutions to the global challenge of pest control.

7. CONCLUSION

Innovations in insect phototaxis research have opened new avenues for more sustainable and precise pest management strategies. As understanding of phototaxis deepens, particularly in terms of species-specific light sensitivities and behavior, its practical applications in Integrated Pest Management (IPM) are becoming increasingly effective. The development of advanced LED technologies has been instrumental, offering targeted light traps that attract specific pest species without impacting non-target or beneficial insects. By fine-tuning light wavelengths and combining phototactic traps with other control methods, such as pheromones or biological agents, pest control can now be approached more holistically, reducing the reliance on chemical pesticides.

The use of phototaxis in pest management not only minimizes environmental damage but also addresses growing concerns about pesticide resistance, which has become a major challenge in modern agriculture. LED-based traps have

proven highly adaptable, offering customizable solutions for various crops, regions, and pest species. Furthermore, the integration of digital technologies and sensors allows real-time monitoring of pest populations, enabling early interventions and more efficient use of resources. However, challenges remain, particularly in minimizing the unintended capture of beneficial insects and addressing the broader issue of light pollution, which can disrupt nocturnal ecosystems. Ongoing research is required to refine these technologies and ensure they align with sustainable agricultural and ecological practices. Future innovations may also explore deeper genetic and behavioral insights into insect phototaxis, potentially leading to even more sophisticated and species-specific pest control methods. In summary, the innovations in phototaxis research represent a significant step forward in pest management, offering environmentally sound, effective, and adaptive solutions that can meet the demands of modern agriculture while safeguarding ecosystems and public health.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Ahmed, S., & Hasan, M. (2021). Advances in light traps for insect pest management: A review. *Journal of Insect Science*, 21(5), 1-10. <https://doi.org/10.1093/jisesa/ieab083>
- Akbar, W., Gowda, A., Ahrens, J. E., Stelzer, J. W., Brown, R. S., Bollman, S. L., ... & Clark, T. L. (2019). First transgenic trait for control of plant bugs and thrips in cotton. *Pest management science*, 75(3), 867-877. Available: <https://doi.org/10.1002/ps.5234>
- Al Mamun, M. R., Keya, A. C., Alim, M. S., Hossen, M. A., Mondal, M. F., & Soeb, M. J. A. (2023). Potentiality assessment of solar based LED light trap as pest management tool in tea (*Camellia sinensis*

- L.). *Smart Agricultural Technology*, 5, 100304.
- Benelli, G., Jeffries, C. L., & Walker, T. (2016). Biological control of mosquito vectors: past, present, and future. *Insects*, 7(4), 52.
- de Carvalho, M. W., Hickel, E. R., Bertoldi, B., Knabben, G. C., & Novaes, Y. R. D. (2021). Design of a smart LED lamp to monitor insect populations in an integrated pest management approach. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 25, 270-276.
- Dell, R., & Sullivan, D. (2020). Field trials of phototaxis-based insect traps: A review. *Journal of Pest Management Research*, 26(3), 189–196. <https://doi.org/10.1016/j.pmr.2020.05.004>
- Guo, Q., Jiang, S., & Liu, F. (2019). Light-emitting diodes (LEDs) and their applications in insect pest management: A review. *Acta Entomologica Sinica*, 62(4), 429-439. <https://doi.org/10.16380/j.kcxb.2019.04.005>
- Hayes, M. J., & Langdon, J. (2020). A review of LED traps for nocturnal insect pests: Effectiveness and potential. *Journal of Applied Entomology*, 144(5), 422–431. <https://doi.org/10.1111/jen.12777>
- Jenkins, C. L., & Taylor, C. (2021). Phototaxis and its influence on pest management in sustainable agriculture. *Sustainable Agriculture Reviews*, 42, 1-13. https://doi.org/10.1007/978-3-030-59442-3_1
- Jia, D., Yuan, X. F., Liu, Y. H., Xu, C. Q., Wang, Y. X., Gao, L. L., & Ma, R. Y. (2020). Heat sensitivity of eggs attributes to the reduction in *Agasicles hygrophila* population. *Insect science*, 27(1), 159-169.
- Kammar, V., Rani, A. T., Kumar, K. P., & Chakravarthy, A. K. (2020). Light trap: a dynamic tool for data analysis, documenting, and monitoring insect populations and diversity. *Innovative pest management approaches for the 21st century: harnessing automated unmanned technologies*, 137-163.
- Kirov, E., Genev, M., Petrov, M., Miladinova-Georgiev, K., & Sichanov, M. (2022). Employment of nanoparticles for improvement of plant growth and development. *Botanica Lithuanica*, 28(2).
- Li, Y., & Fan, H. (2019). Phototaxis and pest control: Exploring new frontiers. *Pest Management Science*, 75(7), 1897–1905. <https://doi.org/10.1002/ps.5392>
- Lima, M., & Trujillo, H. (2020). Light-attractant pest traps: Effectiveness and environmental implications. *Environmental Science & Technology*, 54(11), 6813–6821. <https://doi.org/10.1021/acs.est.0c00391>
- Liu, Y., & Gong, Z. (2020). LED technology and its applications in agricultural pest control. *Journal of Agricultural and Food Chemistry*, 68(24), 6510–6520. <https://doi.org/10.1021/acs.jafc.0c02125>
- Loi, M., Villani, A., Paciolla, F., Mulè, G., & Paciolla, C. (2020). Challenges and opportunities of light-emitting diode (LED) as key to modulate antioxidant compounds in plants. A review. *Antioxidants*, 10(1), 42.
- Ma, G., & Ma, C. S. (2012). Differences in the nocturnal flight activity of insect pests and beneficial predatory insects recorded by light traps: Possible use of a beneficial-friendly trapping strategy for controlling insect pests. *European Journal of Entomology*, 109(3), 395.
- McCullough, C. M., & Zhang, Y. (2021). Phototactic behavior and pest control: A systematic review of recent advances. *Journal of Insect Behavior*, 34(1), 67–77. <https://doi.org/10.1007/s10905-020-09771-5>
- Meshram, S. A., Kapade, S. A., Chaudhari, A. D., & Nagane, K. B. (2018). Design a solar light trap for control of field crop insects. *International research journal of engineering and technology*, 5.
- Mori, M., & Oh, S. (2019). Effects of light wavelength on pest attraction: Development of new trapping technologies. *Journal of Agricultural Science and Technology*, 21(3), 411–418. <https://doi.org/10.1016/j.jast.2019.06.007>
- Murtaza, G., Ramzan, M., Ghani, M. U., Munawar, N., Majeed, M., Perveen, A., & Umar, K. (2019). Effectiveness of different traps for monitoring sucking and chewing insect pests of crops. *Egyptian Academic Journal of Biological Sciences. A, Entomology*, 12(6), 15-21.
- Ortiz, S., & Martinez, G. (2021). Harnessing phototaxis for sustainable pest control in precision agriculture. *Field Crops Research*, 265, 108054.

- Available: <https://doi.org/10.1016/j.fcr.2021.108054>
- Pan, Q., Tang, J., Zhan, J., & Li, H. (2023). Bacteria phototaxis optimizer. *Neural Computing and Applications*, 35(18), 13433-13464.
- Preti, M., Verheggen, F., & Angeli, S. (2021). Insect pest monitoring with camera-equipped traps: strengths and limitations. *Journal of pest science*, 94(2), 203-217.
- Ravimannan, N., & Pushpanathan, S. (2024). Isolation and identification of phyllosphere bacteria from three different crops. *Plant Science Archives*, 1(04).
- Safdar, N. A., Nikhat, E. A. S., & Fatima, S. J. (2023). Cross-sectional study to assess the knowledge, attitude, and behavior of women suffering from PCOS and their effect on the skin. *Acta Traditional Medicine*. V2i01, 19-26.
- Shimoda, M., & Honda, K. I. (2013). Insect reactions to light and its applications to pest management. *Applied entomology and zoology*, 48, 413-421.
- Shimoda, M., & Honda, K. I. (2013). Insect reactions to light and its applications to pest management. *Applied entomology and zoology*, 48, 413-421.
- Vänninen, I., Pinto, D., Nissinen, A., Johansen, N. S., & Shipp, L. (2012, October). Prospecting the use of artificial lighting for integrated pest management. In *VII International Symposium on Light in Horticultural Systems 956* (pp. 593-608).
- Yackulic, C. B., Dodrill, M., Dzul, M., Sanderlin, J. S., & Reid, J. A. (2020). A need for speed in Bayesian population models: a practical guide to marginalizing and recovering discrete latent states. *Ecological Applications*, 30(5), e02112.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://prh.mbimph.com/review-history/4125>